



## Generalizations of a theorem of Herman and a new proof of the simplicity of $\text{Diff}_c^\infty(M)_0$

TOMASZ RYBICKI

Let  $M$  be a smooth manifold of dimension  $n$ . By  $\text{Diff}_c^\infty(M)$  we will denote the group of compactly supported diffeomorphisms of  $M$ . We shall consider a Lie group structure on  $\text{Diff}_c^\infty(M)$  in the sense of the convenient setting of Kriegl and Michor [10]. In particular, we assume that  $\text{Diff}_c^\infty(M)$  is endowed with the  $c^\infty$ -topology [10, Section 4], i.e. the final topology with respect to all smooth curves. For compact  $M$  the  $c^\infty$ -topology on  $\text{Diff}_c^\infty(M)$  coincides with the Whitney  $C^\infty$ -topology, cf. [10, Theorem 4.11(1)]. In general the  $c^\infty$ -topology on  $\text{Diff}_c^\infty(M)$  is strictly finer than the one induced from the Whitney  $C^\infty$ -topology, cf. [10, Section 4.26]. The latter coincides with the inductive limit topology  $\lim_K \text{Diff}_K^\infty(M)$  where  $K$  runs through all compact subsets of  $M$ .

Given smooth complete vector fields  $X_1, \dots, X_N$  on  $M$ , we consider the map

$$(1) \quad \begin{aligned} K: \text{Diff}_c^\infty(M)^N &\rightarrow \text{Diff}_c^\infty(M), \\ K(g_1, \dots, g_N) &:= [g_1, \exp(X_1)] \circ \dots \circ [g_N, \exp(X_N)]. \end{aligned}$$

Here  $\exp(X)$  denotes the flow of a complete vector field  $X$  at time 1, and  $[k, h] := k \circ h \circ k^{-1} \circ h^{-1}$  denotes the commutator of two diffeomorphisms  $k$  and  $h$ . It is readily checked that  $K$  is smooth. Indeed, one only has to observe that  $K$  maps smooth curves to smooth curves, cf. [10, Section 27.2]. Clearly  $K(\text{id}, \dots, \text{id}) = \text{id}$ .

A smooth local right inverse at the identity for  $K$  consists of an open neighborhood  $\mathcal{U}$  of the identity in  $\text{Diff}_c^\infty(M)$  together with a smooth map

$$\sigma = (\sigma_1, \dots, \sigma_N): \mathcal{U} \rightarrow \text{Diff}_c^\infty(M)^N$$

so that  $\sigma(\text{id}) = (\text{id}, \dots, \text{id})$  and  $K \circ \sigma = \text{id}_{\mathcal{U}}$ . More explicitly, we require that each  $\sigma_i: \mathcal{U} \rightarrow \text{Diff}_c^\infty(M)$  is smooth with  $\sigma_i(\text{id}) = \text{id}$  and, for all  $g \in \mathcal{U}$ ,

$$g = [\sigma_1(g), \exp(X_1)] \circ \dots \circ [\sigma_N(g), \exp(X_N)].$$

The aim of this talk is to present the following two results which generalize a well-known theorem of Herman for  $M$  being the torus [8, 9].

---

Key words and phrases. diffeomorphism group; perfect group; simple group; fragmentation; convenient calculus; foliation.

Joint with Stefan Haller and Josef Teichmann.

© 2013 Tomasz Rybicki

**Theorem 1.** *Suppose  $M$  is a smooth manifold of dimension  $n \geq 2$ . Then there exist four smooth complete vector fields  $X_1, \dots, X_4$  on  $M$  so that the map  $K$ , see (1), admits a smooth local right inverse at the identity,  $N = 4$ . Moreover, the vector fields  $X_i$  may be chosen arbitrarily close to zero with respect to the strong Whitney  $C^0$ -topology. If  $M$  admits a proper (circle valued) Morse function whose critical points all have index 0 or  $n$ , then the same statement remains true with three vector fields.*

Particularly, on the manifolds  $M = \mathbb{R}^n, S^n, T^n$ ,  $n \geq 2$ , or the total space of a compact smooth fiber bundle  $M \rightarrow S^1$ , three commutators are sufficient. At the expense of more commutators, it is possible to gain further control on the vector fields. More precisely, we have:

**Theorem 2.** *Suppose  $M$  is a smooth manifold of dimension  $n \geq 2$  and set  $N := 6(n + 1)$ . Then there exist smooth complete vector fields  $X_1, \dots, X_N$  on  $M$  so that the map  $K$ , see (1), admits a smooth local right inverse at the identity. Moreover, the vector fields  $X_i$  may be chosen arbitrarily close to zero with respect to the strong Whitney  $C^\infty$ -topology.*

Either of the two theorems implies that  $\text{Diff}_c^\infty(M)_o$ , the connected component of the identity, is a perfect group, provided  $M$  is not  $\mathbb{R}$ . Our proof rests on Herman's result similarly as that of [17] (see [2]), but is otherwise elementary and different from Thurston's approach. In fact we only need Herman's result in dimension 1.

The perfectness of  $\text{Diff}_c^\infty(M)_o$  was already proved by Epstein [5] using ideas of Mather [11, 12] who dealt with the  $C^r$ -case,  $1 \leq r < \infty$ ,  $r \neq n + 1$ . The Epstein–Mather proof is based on a sophisticated construction, and uses the Schauder–Tychonov fixed point theorem. The existence of a presentation

$$g = [h_1, k_1] \circ \dots \circ [h_N, k_N]$$

is guaranteed, but without any further control on the factors  $h_i$  and  $k_i$ . Theorem 1 or 2 actually implies that the universal covering of  $\text{Diff}_c^\infty(M)_o$  is a perfect group. This result is known, too, see [17]. Thurston's proof is based on a result of Herman for the torus [8, 9]. Note that the perfectness of  $\text{Diff}_c^\infty(M)_o$  implies that this group is simple, see Epstein [4]. The methods used in [4] are elementary and actually work for a rather large class of homeomorphism groups.

One could believe that the phenomenon of smooth perfectness described in Theorems 1 and 2 would be also true for some classical diffeomorphism groups which are simple, e.g. for the Hamiltonian diffeomorphism group of a closed symplectic manifold [1], or for the contactomorphism group of an arbitrary co-oriented contact manifold [15]. However, the available methods seem to be useless for possible proofs of their smooth per-

fectness. Another open problem related to the above theorems is whether a smooth *global* right inverse at the identity for  $K$  would exist. A possible answer in the affirmative seems to be equally difficult. Consequently, it would be difficult to improve Theorems 1 and 2 as they are in any possible direction.

Another essential and important way to generalize the simplicity theorems for  $\text{Diff}_c^\infty(M)_o$ , where  $1 \leq r \leq \infty$ ,  $r \neq n + 1$ , is to consider the uniform perfectness or, more generally, the boundedness of the groups in question. In particular, we ask if the presentation  $g = [h_1, k_1] \circ \cdots \circ [h_N, k_N]$  is available for all  $g \in \text{Diff}_c^\infty(M)_o$  with  $N$  bounded. This property has been proved in the recent papers by Burago, Ivanov and Polterovich [3], and Tsuboi [18], [19], [20], for a large class of manifolds. For instance,  $N = 10$  was obtained in [3] for any closed three dimensional manifold, and then it was improved in [18] to  $N = 6$  for any closed odd dimensional manifold. It seems that the methods of [3], [18], [19] and [20] combined with our Theorem 2 would give some analogue of Theorem 1, but certainly not with the presentation (1) and the condition on  $X_i$ . Also  $N$  could not be smaller in this way. Another advantage of Theorem 1 is that it is valid for all smooth paracompact manifolds. See also [16] for diffeomorphism groups with no restriction of support.

Let  $T^n := \mathbb{R}^n/\mathbb{Z}^n$  denote the torus. For  $\lambda \in T^n$  we let  $R_\lambda \in \text{Diff}^\infty(T^n)$  denote the corresponding rotation. The main ingredient in the proof of Theorems 1 and 2 is the following result of Herman [9, 8].

**Theorem 3** (Herman). *There exist  $\gamma \in T^n$  so that the smooth map*

$$T^n \times \text{Diff}^\infty(T^n) \rightarrow \text{Diff}^\infty(T^n), \quad (\lambda, g) \mapsto R_\lambda \circ [g, R_\gamma],$$

*admits a smooth local right inverse at the identity. Moreover,  $\gamma$  may be chosen arbitrarily close to the identity in  $T^n$ .*

Herman's result is an application of the Nash–Moser inverse function theorem. When inverting the derivative one is quickly led to solve the linear equation  $Y = X - (R_\gamma)^*X$  for given  $Y \in C^\infty(T^n, \mathbb{R}^n)$ . This is accomplished using Fourier transformation. Here one has to choose  $\gamma$  sufficiently irrational so that tame estimates on the Sobolev norms of  $X$  in terms of the Sobolev norms of  $Y$  can be obtained. The corresponding small denominator problem can be solved due to a number theoretic result of Khintchine.

We shall make use of the following corollary of Herman's result.

**Proposition 1.** *There exist smooth vector fields  $X_1, X_2, X_3$  on  $T^n$  so that the smooth map  $\text{Diff}^\infty(T^n)^3 \rightarrow \text{Diff}^\infty(T^n)$ ,*

$$(g_1, g_2, g_3) \mapsto [g_1, \exp(X_1)] \circ [g_2, \exp(X_2)] \circ [g_3, \exp(X_3)],$$

admits a smooth local right inverse at the identity. Moreover, the vector fields  $X_i$  may be chosen arbitrarily close to zero with respect to the Whitney  $C^\infty$ -topology.

The following lemma leads to a decomposition of a diffeomorphism into factors which are leaf preserving. If  $\mathcal{F}$  is a smooth foliation of  $M$  we let  $\text{Diff}_c^\infty(M; \mathcal{F})$  denote the group of compactly supported diffeomorphisms preserving the leaves of  $\mathcal{F}$ . This is a regular Lie group modelled on the convenient vector space of compactly supported smooth vector fields tangential to  $\mathcal{F}$ .

**Lemma 1.** *Suppose  $M_1$  and  $M_2$  are two finite dimensional smooth manifolds and set  $M := M_1 \times M_2$ . Let  $\mathcal{F}_1$  and  $\mathcal{F}_2$  denote the foliations with leaves  $M_1 \times \{\text{pt}\}$  and  $\{\text{pt}\} \times M_2$  on  $M$ , respectively. Then the smooth map*

$$F: \text{Diff}_c^\infty(M; \mathcal{F}_1) \times \text{Diff}_c^\infty(M; \mathcal{F}_2) \rightarrow \text{Diff}_c^\infty(M), \quad F(g_1, g_2) := g_1 \circ g_2,$$

*is a local diffeomorphism at the identity.*

Now we need a version of the exponential law.

**Lemma 2.** *Suppose  $B$  and  $T$  are finite dimensional smooth manifolds, assume  $T$  compact, and let  $\mathcal{F}$  denote the foliation with leaves  $\{\text{pt}\} \times T$  on  $B \times T$ . Then the canonical bijection*

$$C_c^\infty(B, \text{Diff}^\infty(T)) \xrightarrow{\cong} \text{Diff}_c^\infty(B \times T; \mathcal{F})$$

*is an isomorphism of regular Lie groups.*

Another ingredient of the proof is a smooth fragmentation of diffeomorphisms.

Suppose  $U \subseteq M$  is an open subset. Every compactly supported diffeomorphism of  $U$  can be regarded as a compactly supported diffeomorphism of  $M$  by extending it identically outside  $U$ . The resulting injective homomorphism  $\text{Diff}_c^\infty(U) \rightarrow \text{Diff}_c^\infty(M)$  is clearly smooth. Note, however, that a curve in  $\text{Diff}_c^\infty(U)$ , which is smooth when considered as a curve in  $\text{Diff}_c^\infty(M)$ , need not be smooth as a curve into  $\text{Diff}_c^\infty(U)$ . Nevertheless, if there exists a closed subset  $A$  of  $M$  with  $A \subseteq U$  and if the curve has support contained in  $A$ , then one can conclude that the curve is also smooth in  $\text{Diff}_c^\infty(U)$ .

**Proposition 2 (Fragmentation).** *Let  $M$  be a smooth manifold of dimension  $n$ , and suppose  $U_1, \dots, U_k$  is an open covering of  $M$ , ie.  $M = U_1 \cup \dots \cup U_k$ . Then the smooth map*

$$P: \text{Diff}_c^\infty(U_1) \times \dots \times \text{Diff}_c^\infty(U_k) \rightarrow \text{Diff}_c^\infty(M), \quad P(g_1, \dots, g_k) := g_1 \circ \dots \circ g_k,$$

admits a smooth local right inverse at the identity.

Proceeding as in [3] permits to reduce the number of commutators considerably, see also [18] and [19].

**Proposition 3.** *Let  $M$  be a smooth manifold of dimension  $n \geq 2$  and put  $N = 6(n + 1)$ . Moreover, let  $U$  an open subset of  $M$  and suppose  $\phi \in \text{Diff}^\infty(M)$ , not necessarily with compact support, such that the closures of the subsets*

$$U, \phi(U), \phi^2(U), \dots, \phi^N(U)$$

*are mutually disjoint. Then there exists a smooth complete vector field  $X$  on  $M$ , a  $C^\infty$ -open neighborhood  $\mathcal{U}$  of the identity in  $\text{Diff}_c^\infty(U)$ , and smooth maps  $\varrho_1, \varrho_2: \mathcal{U} \rightarrow \text{Diff}_c^\infty(M)$  so that  $\varrho_1(\text{id}) = \varrho_2(\text{id}) = \text{id}$  and, for all  $g \in \mathcal{U}$ ,*

$$g = [\varrho_1(g), \phi] \circ [\varrho_2(g), \exp(X)].$$

*Moreover, the vector field  $X$  may be chosen arbitrarily close to zero in the strong Whitney  $C^\infty$ -topology on  $M$ .*

Now, by applying the Morse theory ([13], [14]) we get

**Lemma 3.** *Let  $M$  be a smooth manifold of dimension  $n$ . Then there exists an open covering  $M = U_1 \cup U_2 \cup U_3$  and smooth complete vector fields  $X_1, X_2, X_3$  on  $M$  so that  $\exp(X_1)(U_1) \subseteq U_2$ ,  $\exp(X_2)(U_2) \subseteq U_3$ , and such that the closures of the sets*

$$U_3, \exp(X_3)(U_3), \exp(X_3)^2(U_3), \dots$$

*are mutually disjoint. Moreover, the vector fields  $X_1, X_2, X_3$  may be chosen arbitrarily close to zero with respect to the strong Whitney  $C^0$ -topology. If  $M$  admits a proper (circle valued) Morse function whose critical points all have index 0 or  $n$ , then we may, moreover, choose  $U_1 = \emptyset$  and  $X_1 = 0$ .*

Theorem 1 is then a consequence of Lemma 3.

## REFERENCES

- [1] A. Banyaga, *Sur la structure du groupe des difféomorphismes qui préservent une forme symplectique*, Comment. Math. Helv. 53 (1978), 174-227.
- [2] A. Banyaga, *The structure of classical diffeomorphism groups*, Kluwer Academic Publishers Group, Dordrecht, 1997.
- [3] D. Burago, S. Ivanov and L. Polterovich, *Conjugation-invariant norms on groups of geometric origin*, Groups of diffeomorphisms, 221–250, Adv. Stud. Pure Math. 52, Math. Soc. Japan, Tokyo, 2008.

- [4] D.B.A. Epstein, *The simplicity of certain groups of homeomorphisms*, *Compositio Math.* **22**(1970), 165–173.
- [5] D.B.A. Epstein, *Commutators of  $C^\infty$ -diffeomorphisms. Appendix to: “A curious remark concerning the geometric transfer map” by John N. Mather*, *Comment. Math. Helv.* **59**(1984), 111–122.
- [6] S. Haller, T. Rybicki and J. Teichmann, *Smooth perfectness for the group of diffeomorphisms*, arXiv:math/0409605
- [7] S. Haller and J. Teichmann, *Smooth perfectness through decomposition of diffeomorphisms into fiber preserving ones*, *Ann. Global Anal. Geom.* **23**(2003), 53–63.
- [8] M.R. Herman, *Simplicité du groupe des difféomorphismes de classe  $C^\infty$ , isotopes à l’identité, du tore de dimension  $n$* , *C. R. Acad. Sci. Paris Sr. A* **273**(1971), 232–234.
- [9] M.R. Herman, *Sur le groupe des difféomorphismes du tore*, *Ann. Inst. Fourier (Grenoble)* **23**(1973), 75–86.
- [10] A. Kriegl and P.W. Michor, *The convenient setting of global analysis*. *Mathematical Surveys and Monographs* **53**, American Mathematical Society, 1997.
- [11] J.N. Mather, *Commutators of diffeomorphisms*, *Comment. Math. Helv.* **49**(1974), 512–528.
- [12] J.N. Mather, *Commutators of diffeomorphisms. II*, *Comment. Math. Helv.* **50**(1975), 33–40.
- [13] J. Milnor, *Morse theory*. *Annals of Mathematics Studies* **51**, Princeton University Press, Princeton, N.J. 1963.
- [14] S.P. Novikov, *Multivalued functions and functionals. An analogue of the Morse theory*, *Dokl. Akad. Nauk SSSR* **260**(1981), 31–35. English translation: *Soviet Math. Dokl.* **24**(1981), 222–226(1982).
- [15] T. Rybicki, *Commutators of contactomorphisms*, *Adv. Math.* **225**(2010), 3291–3326.
- [16] T. Rybicki, *Boundedness of certain automorphism groups of an open manifold*, *Geometriae Dedicata* **151,1**(2011), 175–186.
- [17] W. Thurston, *Foliations and groups of diffeomorphisms*, *Bull. Amer. Math. Soc.* **80**(1974), 304–307.
- [18] T. Tsuboi, *On the uniform perfectness of diffeomorphism groups*. *Groups of diffeomorphisms*, 505–524, *Adv. Stud. Pure Math.* **52**, Math. Soc. Japan, Tokyo, 2008.
- [19] T. Tsuboi, *On the uniform simplicity of diffeomorphism groups*. *Differential geometry*, 43–55, World Sci. Publ., Hackensack, NJ, 2009.
- [20] T. Tsuboi, *On the uniform perfectness of the groups of diffeomorphisms of even-dimensional manifolds*, *Comment. Math. Helv.* **87,1**(2012), 141–185.

Faculty of applied mathematics  
AGH University of Science and Technology  
Al. Mickiewicza 30, 30-059 Krakow, Poland  
E-mail: tomasz@agh.edu.pl